

POSSIBLE BEHAVIOURS OF THE REFLECTION ORDERING OF STATIONARY SETS

JIRÍ WITZANY

The Pennsylvania State University and Charles University (Prague)

February 3, 1994

ABSTRACT. If S, T are stationary subsets of a regular uncountable cardinal κ , we say that S reflects fully in T , $S < T$, if for almost all $\alpha \in T$ (except a nonstationary set) $S \cap \alpha$ is stationary in α . This relation is known to be a well-founded partial ordering. We say that a given poset P is realized by the reflection ordering if there is a maximal antichain $\langle X_p; p \in P \rangle$ of stationary subsets of $\text{Reg}(\kappa)$ so that

$$\forall p, q \in P \forall S \subseteq X_p, T \subseteq X_q \text{ stationary} : (S < T \leftrightarrow p <_P q).$$

We prove that if $V = L[\vec{U}]$, $\mathcal{M}(\kappa) = \kappa^{++}$, and P is an arbitrary well-founded poset of cardinality $\leq \kappa^+$ then there is a generic extension where P is realized by the reflection ordering on κ .

1. Introduction.

If S is a stationary subset of a regular uncountable cardinal κ then *the trace of S* is the set

$$\text{Tr}(S) = \{\alpha < \kappa; S \cap \alpha \text{ is stationary in } \alpha\}$$

and we say that S *reflects at* $\alpha \in \text{Tr}(S)$. If S and T are both stationary, we define

$$S < T \text{ if for almost all } \alpha \in T, \alpha \in \text{Tr}(S)$$

and say that S *reflects fully in T* . (Throughout the paper, “for almost all” means “except for a nonstationary set of points”). It can be proved that this relation is a

1991 *Mathematics Subject Classification.* 03E35, 03E55.

Key words and phrases. Stationary sets, reflection, measurable cardinals, repeat points.

I want to express my gratitude to T.Jech and also to J.Zapletal for many valuable discussions and remarks on the subject.

well-founded partial ordering (see [JW93] or [J84]). The order $o(S)$ of a stationary set of regular cardinals is defined as the rank of S in the relation $<$:

$$o(S) = \sup\{o(T) + 1; T \subseteq \text{Reg}(\kappa) \text{ is stationary and } T < S\}.$$

For a stationary set T such that $T \cap \text{Sing}(\kappa)$ is stationary define $o(T) = -1$. The order of κ is then defined as

$$o(\kappa) = \sup\{o(S) + 1; S \subseteq \kappa \text{ is stationary}\}.$$

Note that if $\text{Tr}(S)$, where $S \subseteq \text{Reg}(\kappa)$, is stationary then $o(S) < o(\text{Tr}(S))$ as $S < \text{Tr}(S)$. It follows from [J84] that the order $o(\kappa)$ provides a natural generalization of the Mahlo hierarchy: κ is exactly $o(\kappa)$ -Mahlo if $o(\kappa) < \kappa^+$ and greatly Mahlo if $o(\kappa) \geq \kappa^+$.

Let P be a well-founded poset, we say that the reflection order $<$ realizes P if there is a maximal antichain $\langle X_p; p \in P \rangle$ of stationary subsets of $\text{Reg}(\kappa)$ so that

$$\forall p, q \in P \forall S \subseteq X_p, T \subseteq X_q \text{ stationary} : (S < T \leftrightarrow p <_P q).$$

If $|P| \leq \kappa$ then it follows that for any stationary $S \subseteq \text{Reg}(\kappa)$

$$\text{Tr}(S) = \sum \{X_p; \exists q \in P : X_q \cap S \text{ is stationary and } q <_P p\}$$

in the Boolean algebra $\mathcal{P}(\text{Reg}(\kappa))/\text{NS}$. Moreover for any $S \subseteq X_p$ stationary $o(S) = o(X_p) = o_P(p)$ where $o_P(p)$ is the rank of p in P and $o(\kappa) = o(P) = \sup\{o_P(p) + 1; p \in P\}$.

In case P is linearly ordered and $S < \text{Reg}(\kappa)$ for any stationary $S \subseteq \text{Sing}(\kappa)$ we say that the Axiom of Full Reflection holds at κ [JS93], the sets X_p are then the canonical stationary sets (see [J84]). [JS93] proves realization of well-orderings of length $\leq \kappa^+$, [JW93] even of length $> \kappa^+$.

Note that only posets of cardinality $\leq \kappa^+$ can be realized as we have only κ^+ subsets of κ (assuming GCH throughout the paper).

A sequence $S = \langle S_\lambda; \lambda \leq \kappa \rangle$ is called a closed system of measures (see [Mi83]) if

$$\forall \lambda \leq \kappa \forall W \in S_\lambda : (j_W S)(\lambda) \subseteq S_\lambda$$

where j_W is the canonical embedding $j_W : V \rightarrow V^\lambda/W$. For $U, W \in S_\lambda$ ($\lambda \leq \kappa$) define $U \triangleleft_S W$ if $U \in (j_W S)(\lambda)$. It is easy to prove that $\triangleleft_S$ is transitive. The standard Mitchell ordering extends $\triangleleft_S$, consequently $\triangleleft_S$ must be well-founded. Let $o_S(U)$ denote the rank of U in this ordering. If $\triangleleft_S$ is linear (on all S_λ) then the system is called a coherent sequence of measures (see [Mi80], [JW93]). We say that the measures in S_κ are separable if there are sets $\langle X_U; U \in S_\kappa \rangle$ so that

$$\forall U, W \in S_\kappa : X_U \in W \text{ iff } U = W.$$

If $S = \langle S_\lambda; \lambda \leq \kappa \rangle$ is a closed system of measures then we say that a $U \in S_\kappa$ is a repeat point if

$$\forall X \in U \exists W \triangleleft_S U : X \in W$$

i.e. U is not separable from its predecessors. It has been proved in [JW93] that if S is a coherent sequence and there are no repeat points in S_κ then the measures in S_κ are separable.

In section 2 the construction of [JW93] is generalized to show the following

Theorem 1. *If $S = \langle S_\lambda; \lambda \leq \kappa \rangle$ is a closed system of separable measures then there is a generic extension $V[G]$ preserving cardinalities, cofinalities, and GCH where the reflection ordering of stationary subsets of $\text{Reg}(\kappa)$ realizes the poset $(S_\kappa, \triangleleft_S)$ (as computed in V).*

Section 3 analyzes the question what well-founded posets are representable by $(S_\kappa, \triangleleft_S)$ and when the measures in S_κ are separable. It turns that closed systems of measures can be easily constructed using a Laver's function on κ that exists in $L[\vec{U}]$ if $\mathfrak{o}^\mathcal{U}(\kappa) = \kappa^{++}$.

Theorem 2. *Assume that $V = L[\vec{U}]$, $\mathfrak{o}^\mathcal{U}(\kappa) = \kappa^{++}$, and that P is a well-founded poset of cardinality $\leq \kappa^+$. Then there is a closed system of measures $\langle S_\lambda; \lambda \leq \kappa \rangle$ such that $P \cong (S_\kappa, \triangleleft_S)$ and the measures in S_κ are separable.*

Corollary. *If $V = L[\vec{U}]$, $\mathfrak{o}^\mathcal{U}(\kappa) = \kappa^{++}$, then any well-founded poset of cardinality $\leq \kappa^+$ is realized by the reflection ordering of stationary subsets of $\text{Reg}(\kappa)$ in a generic extension of V preserving cofinalities, cardinalities, and GCH.*

2. The forcing construction.

The construction is analogous to the construction of [JW93]. We will spell out its definition but will not repeat the proofs that are almost literally same as the proofs in [JW93].

Let $S = \langle S_\lambda; \lambda \leq \kappa \rangle$ be a closed system of measures in the ground model V satisfying GCH.

As usual, if P is a forcing notion then $V(P)$ denotes either the Boolean valued model or a generic extension by a P -generic filter over V .

$P_{\kappa+1}$ will be an Easton support iteration of $\langle Q_\lambda; \lambda \leq \kappa \rangle$, Q_λ will be nontrivial only for λ Mahlo. Q_λ (for λ Mahlo) is defined in $V(P_\lambda)$, where P_λ denotes the iteration below λ , as an iteration of length λ^+ with $< \lambda$ -support of forcing notions shooting clubs through certain sets $X \subseteq \lambda$ (we will denote this standard forcing notion $CU(X)$), always with the property that $X \supseteq \text{Sing}(\lambda)$. This condition will guarantee Q_λ to be essentially $< \lambda$ -closed (i.e. for any $\gamma < \lambda$ there is a dense γ -closed subset of Q_λ). Q_λ will also satisfy the λ^+ -chain condition. Consequently P_λ will satisfy λ -c.c. and will have size λ . Cardinalities, cofinalities, and GCH will be preserved, stationary subsets of λ can be made nonstationary only by the forcing at λ , not below λ , and not after the stage λ - after stage λ no subsets of λ are added.

We use the λ^+ -chain condition of Q_λ to get a canonical enumeration of length λ^+ of all Q_λ -names for subsets of λ so that the β th name appears in $V(P_\lambda * Q_\lambda | \beta)$. Moreover for $U \in S_\lambda$ we will define certain filters F_U in $V(P_\lambda * Q_\lambda | \beta)$. Their definition will not be absolute, however the filter F_U will extend the measure U and will increase coherently during the iteration.

Definition. An iteration Q of $\langle CU(B_\alpha); \alpha < \alpha_0 \rangle$ with $< \lambda$ -support and length $\alpha < \lambda^+$ is called an $\tilde{S}$ -iteration (for $\tilde{S} \subseteq S_\lambda$) if for all $\alpha < \alpha_0$

$$V(P_\lambda * Q|\alpha) \models "B_\alpha \in F_U \text{ for any } U \in \tilde{S} \text{ and } \text{Sing}(\lambda) \subseteq B_\alpha."$$

An $(S_\lambda|U \cup \{U\})$ -iteration, where $S_\lambda|U = \{W \in S_\lambda; W \triangleleft_S U\}$, is also called an iteration of order U .

(Note that any S_λ -iteration is also an $S_\lambda|U$ -iteration.)

Q_λ is then defined as an iteration of $\langle CU(B_\alpha); \alpha < \lambda^+ \rangle$ with $< \lambda$ -support and length λ^+ so that every $Q_\lambda|\alpha$ is an S_λ -iteration and all potential names $\dot{X} \subseteq \lambda$ are used cofinally many times in the iteration as some B_α .

Observe that Q_λ can be represented in $V(P_\lambda)$ as a set of sequences of closed bounded subsets of λ in $V(P_\lambda)$ rather than in $V(P_\lambda * Q_\lambda|\alpha)$. Moreover if $\dot{q}$ is a P_λ -name such that $1 \Vdash_{P_\lambda} \dot{q} \in Q_\lambda$ then using the λ -chain condition of P_λ there is a set $A \subseteq \lambda^+$ (in V) of cardinality $< \lambda$ and $\gamma_0 < \lambda$ so that

$$1 \Vdash_{P_\lambda} \text{supp } \dot{q} \subseteq A \text{ and } \forall \alpha \in A : \dot{q}(\alpha) \subseteq \gamma_0 .$$

Consequently, Q_λ can be represented as a set of functions $g : A \times \gamma_0 \rightarrow [P_\lambda]^{<\lambda}$ where $A \subseteq \lambda^+$, $|A| < \lambda$ and $\gamma_0 < \lambda$. In this sense Q_λ has cardinality λ^+ and any $Q_\lambda|\alpha$ has cardinality at most λ .

Definition of filters F_U .

The filters F_U ($U \in S_\lambda$) are defined in $V(P_\lambda * Q)$, where Q is any iteration of order U , by induction so that the following is satisfied:

Proposition 2.1. Let $Q, Q' = Q * R$ be two iterations of order U then

$$F_U^{V(P_\lambda * Q)} = F_U^{V(P_\lambda * Q')} \cap V(P_\lambda * Q) .$$

Moreover $F_U^{V(P_\lambda)} \cap V = U$.

Proposition 2.2. Let $j = j_U$ be the canonical embedding from V into $V^\lambda/U = M$ and Q an iteration of order U . Then j can be lifted to an elementary embedding from a generic extension $V(P_\lambda * Q)$ of V to a generic extension $M(jP_\lambda * jQ)$ of M .

Lemma 2.3. Let $N = V^\lambda/W$ for some $W \triangleright_S U$ and Q be an iteration of order U . Then

$$F_U^{V(P_\lambda * Q)} = F_U^{N(P_\lambda * Q)} .$$

Note that it also means that the definition of F_U relativized to $N(P_\lambda * Q)$ makes sense.

Lemma 2.4. *Let $j = j_U : V \rightarrow M$. Then any $S_\lambda|U$ -iteration Q is an subiteration of $(jP_\lambda)^\lambda$, where $(jP_\lambda)^\lambda$ is the factor of $jP_\lambda = P_\lambda * (jP_\lambda)^\lambda * (jP_\lambda)^{>\lambda}$. Consequently for any G^* jP_λ -generic/ V and any $q \in Q$ there is an $H \in M[G^*]$ Q -generic/ $V[G]$ containing q given by an embedding of Q as a subiteration of $(jP_\lambda)^\lambda$, where $G = G^* \restriction P_\lambda$.*

Definition. *Let j , Q , G^* , G be as in the lemma. Then $\text{Gen}_j(Q, G^*)$ denotes the set of all filters $H \in M[G^*]$ Q -generic/ $V[G]$ given by an embedding of Q as a subiteration of $(jP_\lambda)^\lambda$.*

Lemma 2.5. *Let j be as above, Q an iteration of order U , G^* jP_λ -generic/ V , $H \in \text{Gen}_j(Q, G^*)$. For every $\beta < l(Q)$ let $C_\beta \subset \lambda$ be the club $\cup\{r(\beta); r \in H\}$, and let $[H]^j$ denote the $j(l(Q))$ -sequence given by*

$$[H]^j(\gamma) = \begin{cases} C_\beta \cup \{\lambda\}, & \text{if } \gamma = j(\beta) \\ \emptyset & \text{otherwise.} \end{cases}$$

Then $[H]^j \in jQ/G^$.*

Propositions 2.1 and 2.2 are essential to analyze reflection of stationary sets in the generic extension.

Now let $U \in S_\lambda$ and suppose that the filters $F_{U'}$ have been defined for all $\lambda' < \lambda$ and $U' \in S_{\lambda'}$ and for all $U' \in S_\lambda|U$ so that 2.1-2.5 holds. Moreover let $\alpha < \lambda^+$ and F_U be defined for all iterations of order U and length $< \alpha$ so that 2.4 and 2.5 holds for U and iterations of length $\leq \alpha$. Then we can define F_U for iterations of order U and length α .

Definition. *Let Q be an iteration of order U and length α , $j = j_U : V \rightarrow M$. For a $P_\lambda * Q$ -name $\dot{X}$ of a subset of λ and $p * q \in P_\lambda * Q$ define*

$$p * q \Vdash_{P_\lambda * Q} \dot{X} \in F_\delta$$

if the following holds in V :

$$j(p) \Vdash_{jP_\lambda} \text{“For any } H \in \text{Gen}_j(Q, G^*) \text{ containing } q :$$

$$[H]^j \Vdash_{jQ} \check{\lambda} \in j\dot{X} \text{ .”}$$

The definition says that $p * q \Vdash \dot{X} \in F_\delta$ if $\lambda \in j^*X$ whenever $j^* : V[G * H] \rightarrow M[G^* * H^*]$ is a lifting of j of certain kind and $p * q \in G * H$.

For the proofs of 2.1–2.5 see [JW93], modifications are left to the reader.

Reflection of stationary sets in the generic extension.

Let us analyze the behaviour of the reflection ordering of stationary subsets of λ in the generic extension $V(P_{\lambda+1})$. That is the same as in $V(P_{\kappa+1})$ because no subsets of λ are added after the stage λ . Note that any subset of λ in $V(P_{\lambda+1})$ already appears in $V(P_\lambda * Q_\lambda|\alpha)$ for some $\alpha < \lambda^+$.

Lemma 2.6. $V(P_\lambda * Q_\lambda | \alpha) \models \text{Club}(\lambda) \subseteq F_U$ for any $U \in S_\lambda$.

Proof. See 3.2 in [JW93].

Lemma 2.7. Let $S \subseteq \text{Reg}(\lambda)$ be in $V(P_{\lambda+1})$ then S is stationary iff there is a $U \in S_\lambda$ such that for some $\alpha < \lambda^+$ $S \in V(P_\lambda * Q_\lambda | \alpha)$ and $V(P_\lambda * Q_\lambda | \alpha) \models$ “ S is F_U -positive” (i.e. iff this is true for all α such that $S \in V(P_\lambda * Q_\lambda | \alpha)$).

Proof. Let $S \in V(P_\lambda * Q_\lambda | \alpha)$ and $V(P_\lambda * Q_\lambda | \alpha) \models$ “ S is F_U -positive” and suppose by contradiction that $V(P_{\lambda+1}) \models$ “ S is nonstationary”. Then there is an $\alpha' > \alpha$ such that

$$V(P_\lambda * Q_\lambda | \alpha') \models \text{“}S \text{ is nonstationary”}$$

however proposition 2.1 says that

$$V(P_\lambda * Q_\lambda | \alpha') \models \text{“}S \text{ is } F_U\text{-positive”}$$

and that is a contradiction with lemma 2.6.

On the other hand suppose that $S \in V(P_\lambda * Q_\lambda | \alpha)$ and

$$V(P_\lambda * Q_\lambda | \alpha) \models \text{“}S \text{ is } F_U\text{-thin for any } U \in S_\lambda \text{.”}$$

Then by the definition of Q_λ a club is shot through $\kappa \setminus S$, consequently S is nonstationary in $V(P_{\lambda+1})$. $\square$

Lemma 2.8. Assume that S is in $V(P_\lambda * Q_\lambda | \alpha)$, $S \subseteq \text{Reg}(\lambda)$ and $W \in S_\lambda$. Then $\text{Tr}(S)$ is either F_W -thin or it belongs to F_W . The former is true iff S is F_U -thin for all $U \triangleleft_S W$. Moreover if $S \subseteq \text{Sing}(\lambda)$ is stationary then $\text{Tr}(S) \in F_U$ for all $U \in S_\lambda$.

Proof. Suppose that there is $U \triangleleft_S W$ such that $p * q \Vdash_{P_\lambda * Q_\lambda | \alpha}$ “ $\dot{S}$ is F_U -positive” but $p * q \nVdash_{P_\lambda * Q_\lambda | \alpha}$ “ $\text{Tr}(\dot{S}) \in F_W$ ” for some $W \triangleright_S U$. Denote $j = j_W : V \rightarrow M = V^\lambda / W$. Then there is a filter G^* jP_λ -generic/ V , $G^* \ni jp$, and $H \in \text{Gen}_j(Q_\lambda | \alpha, G^*)$, $H \ni q$, and a filter H^* $j(Q_\lambda | \alpha)$ -generic/ $V[G^*]$, H^* contains $[H]^j$, so that $\lambda \notin j\text{Tr}(\dot{S})_{/G^* * H^*}$. The embedding j is lifted to

$$j^{**} : V[G * H] \rightarrow M[G^* * H^*].$$

$S = \dot{S}_{/G * H}$ is F_U -positive in $V[G * H]$ and

$$j\text{Tr}(\dot{S})_{/G^* * H^*} = j^{**}(\text{Tr}(S)) = \text{Tr}^{M[G^* * H^*]}(j^{**}S).$$

Thus $\lambda \notin \text{Tr}^{M[G^* * H^*]}(j^{**}S)$ which means that

$$M[G^* * H^*] \models \text{“}S \text{ is not stationary in } \lambda\text{”}$$

because $S = j^{**}S \cap \lambda$. Consequently

$$V[G^* * H^*] \models "S \text{ is not stationary in } \lambda."$$

Observe that $j(Q_\lambda|\alpha)$ has a dense subset λ -closed in $M[G^*]$ and thus also in $V[G^*]$. Moreover $jP_\lambda = P_\lambda * (jP_\lambda)^\lambda * R$ where R is essentially λ -closed in $V[G^*|\lambda+1]$. It implies that already $V[G^*|\lambda+1] \models "S \text{ is not stationary in } \lambda."$ Let us now consider the isomorphism $(jP_\lambda)^\lambda \simeq (Q_\lambda|\alpha) * \tilde{Q}$ from the proof of 2.4 giving a filter $H = G^* \restriction (Q_\lambda|\alpha)$, let $\tilde{H} = G^* \restriction \tilde{Q}$. Since every subset of λ in $V[G * H * \tilde{H}]$ is already in some $V[G * H * \tilde{H}|\beta]$ there is a $\beta < \lambda^+$ so that

$$V[G * H * \tilde{H}|\beta] \models "S \text{ is not stationary in } \lambda."$$

But since $(Q_\lambda|\alpha) * (\tilde{Q}|\beta)$ is an $S_\lambda|W$ -iteration and hence an iteration of order U it follows from proposition 2.1 that

$$V[G * H * \tilde{H}|\beta] \models "S \text{ is } F_U\text{-positive}"$$

which contradicts lemma 2.6.

The proof that for any stationary $S \subseteq \text{Sing}(\lambda)$ and $U \in S_\lambda$, $\text{Tr}(S) \in F_U$ is analogous using the following fact instead of proposition 2.1.

Claim. *Stationary subsets of $\text{Sing}(\lambda)$ are preserved by $\emptyset$ -iterations.*

Proof. See 7.38 in [J86] or 3.4 in [JW93].

Now let $S \in V(P_\lambda * Q_\lambda|\alpha)$, $W \in S_\lambda$ and $S \subseteq \text{Reg}(\lambda)$ be F_U -thin for any $U \triangleleft_S W$. We want to prove that $V(P_\lambda * Q_\lambda|\alpha) \models \lambda \setminus \text{Tr}(S) \in F_W$. Let $j = j_W : V \rightarrow M$, then $(jP_\lambda)^\lambda$ is an iteration of length λ^+ such that $(jP_\lambda)^\lambda|\beta$ is always an $S_\lambda|W$ -iteration and every potential name is used cofinally many times. For β large enough $Q_\lambda|\alpha$ is an subiteration of $(jP_\lambda)^\lambda|\beta$ and

$$V(P_\lambda * (jP_\lambda)^\lambda|\beta) \models "\lambda \setminus S \in F_U \text{ for any } U \in S_\lambda|W"$$

because $F_U^{V(P_\lambda * (jP_\lambda)^\lambda|\beta)} \supseteq F_U^{V(P_\lambda * Q_\lambda|\alpha)} \ni \lambda \setminus S$. Consequently a club is shot through $\lambda \setminus S$ in the iteration $(jP_\lambda)^\lambda$. It implies that

$$V[G^*] \models "S \subseteq \lambda \text{ is nonstationary}"$$

where G^* is any jP_λ -generic/ V and thus also for any $H \in \text{Gen}_j(Q_\lambda, G^*)$ and any $H^* \ni [H]^j$ $j(Q_\lambda|\alpha)$ -generic/ $V[G^*]$

$$V[G^* * H^*] \models \lambda \in j^*(\lambda \setminus \text{Tr}(S)).$$

That proves $V(P_\lambda * Q_\lambda|\alpha) \models \lambda \setminus \text{Tr}(S) \in F_W$. $\square$

The filters F_U are not defined in $V(P_\lambda * Q_\lambda)$, however we can define $\tilde{F}_U = \bigcup_{\alpha < \lambda^+} F_U^{V(P_\lambda * Q_\lambda | \alpha)}$. It follows from the lemmas that a set $S \subseteq \text{Reg}(\lambda)$ in $V(P_{\lambda+1})$ is stationary iff it is $\tilde{F}_U$ -stationary for some U . Moreover for any U either $\text{Tr}(S) \in \tilde{F}_U$ or $\text{Tr}(S)$ is $\tilde{F}_U$ -thin. The former is true iff there is a $W \triangleleft_S U$ such that S is $\tilde{F}_W$ -positive. Besides for $S \subseteq \text{Sing}(\lambda)$ stationary $\text{Reg}(\lambda) \subseteq \text{Tr}(S) \pmod{\text{NS}}$. If the measures $U \in S_\lambda$ are separated by sets $X_U \subseteq \text{Reg}(\lambda)$ it follows that $\langle X_U; U \in S_\lambda \rangle$ forms a maximal antichain of stationary subsets of $\text{Reg}(\lambda)$ in $V(P_{\lambda+1})$ and the reflection ordering of stationary subsets of $\text{Reg}(\lambda)$ realizes the poset $(S_\lambda, \triangleleft_S)$. Actually $\tilde{F}_U = C[X_U] = \{Y \subseteq \kappa; \exists C \text{ a club} : C \cap X_U \subseteq Y\}$. That proves theorem 1.

3. Closed systems of measures.

In this section we construct closed systems of separable measures isomorphic to a given well-founded poset.

Lemma 3.1. *Let $S = \langle S_\lambda; \lambda \leq \kappa \rangle$ be a system of measures such that*

$$\forall U, W \in S_\kappa : (j_U S)(\kappa) \subseteq S_\kappa \text{ and } (W \in (j_U S)(\kappa) \Rightarrow (j_W S)(\kappa) \subseteq (j_U S)(\kappa)).$$

Then there is a closed subsystem of measures $\tilde{S} = \langle \tilde{S}_\lambda; \lambda \leq \kappa \rangle$ such that $S_\kappa = \tilde{S}_\kappa$ and $\forall U, V \in S_\kappa : U \triangleleft_S V \leftrightarrow U \triangleleft_{\tilde{S}} V$.

Proof. Define $\tilde{S}_\lambda$ by induction on $\lambda \leq \kappa$. Suppose $\tilde{S}_{\lambda'}$ has been defined for $\lambda' < \lambda$. If $\forall U \in S_\lambda : (j_U(\tilde{S} \upharpoonright \lambda))(\lambda) \subseteq S_\lambda$ then put $\tilde{S}_\lambda = S_\lambda$ otherwise $\tilde{S}_\lambda = \emptyset$. That defines $\tilde{S}_\lambda \subseteq S_\lambda$ for all $\lambda \leq \kappa$, obviously $\tilde{S}_\kappa = S_\kappa$ by the assumption of the lemma. Finally let us prove that for any $U \in S_\kappa$

$$\{\lambda < \kappa; \tilde{S}_\lambda = S_\lambda\} \in U$$

and consequently $(j_U S)(\kappa) = (j_U \tilde{S})(\kappa)$. It is enough to prove that the following set is in U

$$A = \{\lambda < \kappa; \forall W \in S_\lambda : (j_W(S \upharpoonright \lambda))(\lambda) \subseteq S_\lambda\}.$$

That is true iff $\kappa \in j_U A$, iff

$$\forall W \in (j_U S)(\kappa) : (j_W^{M_U}(S \upharpoonright \kappa))(\kappa) \subseteq (j_U S)(\kappa)$$

where $j_U : V \rightarrow V^\kappa/U = M_U$ and $j_W^{M_U} : M_U \rightarrow M_U^\kappa/W$. For any $W \in (j_U S)(\kappa) \subseteq S_\kappa$ the embedding $j_W^{M_U} = j_W \upharpoonright M_U$, thus we have $(j_W^{M_U}(S \upharpoonright \kappa))(\kappa) = (j_W S)(\kappa) \subseteq (j_U S)(\kappa)$. $\square$

As a corollary we can prove

Proposition 3.2. *Suppose P is a well-founded poset of cardinality $\leq \kappa$ and there are measures U_p ($p \in P$) on κ so that $U_p \triangleleft U_q$ whenever $p <_P q$. Then there is a*

closed system of separable measures $S = \langle S_\lambda; \lambda \leq \kappa \rangle$ such that $S_\kappa = \{U_p; p \in P\}$ and

$$p <_P q \leftrightarrow U_p \triangleleft_S U_q.$$

Proof. Since $|P| \leq \kappa$ we can find disjoint sets $X_p \in U_p$ ($p \in P$) separating the measures. For $p \in P$ the set $\{U_q; q <_P p\}$ is in the ultraproduct V^κ/U_p because all U_q ($q <_P p$) are and the number of them is at most κ . Let $\langle S_\lambda; \lambda \in X_p \rangle$ be a sequence of sets of measures over λ such that

$$[S_\lambda]_{U_p}^{\lambda \in X_p} = \{U_q; q <_P p\}.$$

Glue together all those sequences into $\langle S_\lambda; \lambda \leq \kappa \rangle$ (put $S_\lambda = \emptyset$ if $\lambda \notin \bigcup_{p \in P} X_p$). Then for any $p \in P$

$$(j_{U_p} S)(\kappa) = \{U_q; q <_P p\}.$$

Finally use lemma 3.1 (the assumption is satisfied as $<_P$ is transitive) to get the desired closed system of measures. $\square$

Closed systems of measures can be easily constructed using a Laver's function.

Definition. (cf. [La78]) *We say that a function $f : \kappa \rightarrow V_\kappa$ is a Laver's function on κ if*

$$\forall x \in V_{\kappa+2} \exists U \text{ a measure on } \kappa : (j_U f)(\kappa) = x.$$

It means that the Laver's function serves as a universal function in ${}^\kappa V$ for all $x \in V_{\kappa+2}$. Generalizing [La78] we can prove that if κ is $\mathcal{P}_2\kappa$ -strong then there is a Laver's function on κ . We will show that the existence of a Laver's function on κ is actually equiconsistent with the Mitchell order of κ being κ^{++} .

It will follow that if there is a Laver's function on κ then there is a coherent sequence of measures $\vec{U}$ such that $o^{\mathcal{U}}(\kappa) = \kappa^{++}$, and moreover that *the measures on κ cover $\mathcal{P}(\kappa^+)$* in the following sense:

$$\forall A \in \mathcal{P}(\kappa^+) \exists \alpha < \kappa^{++} : A \in \text{Ult}(V, U_\alpha^\kappa).$$

Let us prove that those two conditions are sufficient for the existence of a Laver's function.

Proposition 3.3. *Let $\vec{U}$ be a coherent sequence of measures such that $o^{\mathcal{U}}(\kappa) = \kappa^{++}$ and the measures $\{U_\alpha^\kappa; \alpha < \kappa^{++}\}$ cover $\mathcal{P}(\kappa^+)$. Then there is a Laver's function on κ .*

Proof. Firstly observe that $\{U_\alpha^\kappa; \alpha < \kappa^{++}\}$ also covers $V_{\kappa+2}$: let $\pi \in \text{Ult}(V, U_0^\kappa)$ be a bijection between κ^+ and $V_{\kappa+1}$. If $x \in V_{\kappa+2}$, put $A = \pi^{-1}[x]$, and find $\alpha < \kappa^{++}$ such that $A \in \text{Ult}(V, U_\alpha^\kappa)$. Since $\pi \in \text{Ult}(V, U_\alpha^\kappa)$, x must be in $\text{Ult}(V, U_\alpha^\kappa)$. It can be assumed without loss of generality that $o^{\mathcal{U}}(\lambda) < \lambda^{++}$ for all $\lambda < \kappa$.

Claim. *There are well-orderings R^λ of $V_{\lambda+2}$ of order type λ^{++} ($\lambda \leq \kappa$) so that for any $\alpha < o^\mathcal{U}(\kappa)$ the well-ordering $j_{U_\alpha^\kappa}(\langle R^\lambda; \lambda < \kappa \rangle)(\kappa)$ is an initial segment of R^κ .*

Proof. Start with arbitrary well-orderings $<_\lambda$ of $V_{\lambda+2}$ of order type λ^{++} ($\lambda \leq \kappa$). Assume $\langle R^\lambda; \lambda < \mu \rangle$ have been defined so that for all $\lambda < \mu$

$$(3.1) \quad \forall \alpha < o^\mathcal{U}(\lambda) : R_\alpha^\lambda = j_{U_\alpha^\lambda}(\langle R^\lambda; \lambda' < \lambda \rangle)(\lambda) \text{ is an initial segment of } R^\lambda.$$

Let $\beta < o^\mathcal{U}(\mu)$, then by elementarity $R_\beta^\mu = j_{U_\beta^\mu}(\langle R_\lambda; \lambda < \mu \rangle)(\mu)$ is a well-ordering of $V_{\lambda+2} \cap \text{Ult}(V, U_\beta^\mu)$ of order type $\mu^{++\text{Ult}(V, U_\beta^\mu)}$ and

$$\text{Ult}(V, U_\beta^\mu) \models \text{“}\forall \alpha < \beta : R_\alpha^\mu = j_{U_\alpha^\mu}(\langle R_\lambda; \lambda < \mu \rangle)(\mu) \text{ is an initial segment of } R_\beta^\mu\text{.”}$$

Consequently R^μ can be defined as $\cup\{R_\beta^\mu; \beta < o^\mathcal{U}(\mu)\}$ and end-extended by $<_\mu$ on the remaining elements of $V_{\mu+2}$. Then (3.1) is also satisfied for $\lambda = \mu$. If $\mu = \kappa$ then there are no remaining sets in $V_{\kappa+2}$, and $R^\kappa = \cup\{R_\beta^\kappa; \beta < o^\mathcal{U}(\kappa)\}$. $\square$ Claim

Now define f by induction as follows: assume $f(\lambda')$ has been defined for $\lambda' < \lambda$, and let $f(\lambda)$ be the R^λ -least $x \in V_{\lambda+2}$ such that

$$\neg \exists \alpha < o^\mathcal{U}(\lambda) : x = (j_{U_\alpha^\lambda}(f \upharpoonright \lambda))(\lambda).$$

Leave $f(\lambda)$ undefined if there is no such x . Assume there is $x \in V_{\kappa+2}$ such that

$$\neg \exists \alpha < o^\mathcal{U}(\kappa) : x = (j_{U_\alpha^\kappa} f)(\kappa),$$

and let x be the R^κ -least with this property. Let $\beta < o^\mathcal{U}(\kappa)$ be such that $x \in \text{Ult}(V, U_\beta^\kappa)$ and

$$\forall y <_{R^\kappa} x \exists \alpha < \beta : y = (j_{U_\alpha^\kappa} f)(\kappa),$$

this is possible as $o^\mathcal{U}(\kappa) = \kappa^{++}$ and there are at most κ^+ many $y <_{R^\kappa} x$. Then

$$\text{Ult}(V, U_\beta^\kappa) \models \text{“}x \text{ is the } R_\beta^\kappa\text{-least such that } \neg \exists \alpha < o^{j\mathcal{U}}(\kappa) = \beta : x = (j_{U_\alpha^\kappa} f)(\kappa)\text{.”}$$

Consequently $(j_{U_\beta^\kappa} f)(\kappa) = x$ by elementarity - a contradiction. $\square$ 3.3

Let us prove that the assumptions of 3.3 are much weaker than $\mathcal{P}_2\kappa$ -strongness. W.Mitchell proved in [Mi83] that there is an inner model $L[\vec{\mathcal{U}}]$ satisfying GCH such that $\vec{\mathcal{U}}$ is a coherent sequence of measures in $L[\vec{\mathcal{U}}]$, all measures in $L[\vec{\mathcal{U}}]$ are in $\vec{\mathcal{U}}$, and $o^\mathcal{U}(\kappa) = \min(o(\kappa), \kappa^{++L[\vec{\mathcal{U}}]})$ for all ordinals κ .

Proposition 3.4. *If $\mathcal{U}(\kappa) = \kappa^{++}$ in $L[\vec{\mathcal{U}}]$ then the measures on κ cover $\mathcal{P}(\kappa^+)$.*

Proof. Let us firstly suppose that κ is the maximal measurable cardinal in $L[\vec{\mathcal{U}}]$ (it can be easily achieved by cutting the universe at the first measurable above κ and then applying the Mitchell's construction of $L[\vec{\mathcal{U}}]$). Notice that the sequence of measures must be actually represented as $\vec{\mathcal{U}} = \{(\lambda, \alpha, X); X \in U_\alpha^\lambda\}$ in order $L[\vec{\mathcal{U}}]$ makes sense. Let $A \subseteq \kappa^+$, $A \in L_\gamma[\vec{\mathcal{U}}]$, $\gamma > \kappa^+$. Find a model $M \subseteq L_\gamma[\vec{\mathcal{U}}]$ of cardinality κ^+ such that $\kappa^+ + 1 \subseteq M$, $A \in M$, $\mathcal{P}(\kappa) \subseteq M$,

$$\forall \alpha \in M : \alpha < \kappa^{++} \Rightarrow \alpha \subseteq M, \text{ and}$$

$$\langle M, \in, \vec{\mathcal{U}} \cap M \rangle \prec \langle L_\gamma[\vec{\mathcal{U}}], \in, \vec{\mathcal{U}} \rangle.$$

Let π be the transitive collapse of M . Then it is easy to see that $\vec{\mathcal{V}} = \pi(\vec{\mathcal{U}} \cap M) = \vec{\mathcal{U}} \cap M = \vec{\mathcal{U}} \upharpoonright (\kappa, \Theta)$ where $\Theta = M \cap \kappa^{++}$. By 32.7 of [Ka93] $\pi[M] = L_\delta[\vec{\mathcal{U}} \upharpoonright (\kappa, \Theta)]$ for a $\delta < \kappa^{++}$. Consequently

$$A = \pi(A) \in L_\delta[\vec{\mathcal{U}} \upharpoonright (\kappa, \Theta)] \subseteq \text{Ult}(V, U_\Theta^\kappa).$$

The idea of the proof in general situation is due to W. Mitchell (personal communication). Start with $A \subseteq \kappa^+$, let $M, L_\gamma[\vec{\mathcal{U}}], \pi, \vec{\mathcal{V}} = \pi(\vec{\mathcal{U}} \cap M), L_\delta[\vec{\mathcal{V}}] = \pi[M]$ be as above. Then we can prove only that $\vec{\mathcal{V}} \upharpoonright (\kappa + 1) = \vec{\mathcal{U}} \upharpoonright (\kappa, \Theta)$, the measures above κ in $\vec{\mathcal{V}}$ does not have to be same as in $\vec{\mathcal{U}}$. Observe that $L_\delta[\vec{\mathcal{V}}]$ is iterable since it is embedded into iterable $L_\gamma[\vec{\mathcal{U}}]$. By the Mitchell's comparison lemma [Mi83] there are iterations defined in $L[\vec{\mathcal{U}}]$

$$i : L_\delta[\vec{\mathcal{V}}] \rightarrow L_\vartheta[\vec{\mathcal{W}}],$$

$$i' : L[\vec{\mathcal{U}}] \rightarrow L[\vec{\mathcal{W}}']$$

such that either $\vec{\mathcal{W}}$ is an initial segment of $\vec{\mathcal{W}}'$ or $\vec{\mathcal{W}}'$ is an initial segment of $\vec{\mathcal{W}}$. $\vec{\mathcal{W}}$ is an initial segment of $\vec{\mathcal{W}}'$ if $\vec{\mathcal{W}} \cap L_\vartheta[\vec{\mathcal{W}}] \cap L[\vec{\mathcal{W}}'] = \vec{\mathcal{W}}' \upharpoonright \beta \cap L_\vartheta[\vec{\mathcal{W}}] \cap L[\vec{\mathcal{W}}']$ for an ordinal β . Assume this is true, it is then easy to see that $L_\vartheta[\vec{\mathcal{W}}] \subseteq L[\vec{\mathcal{W}}']$. It follows from the proof of the comparison lemma that $\text{crit}(i) > \kappa$, and the iteration i' starts with the ultraprodut by U_Θ^κ and then proceeds with measures above κ . Thus $A \in L_\vartheta[\vec{\mathcal{W}}] \subseteq L[\vec{\mathcal{W}}']$ implies $A \in \text{Ult}(L[\vec{\mathcal{W}}], U_\Theta^\kappa)$. In that case we are done.

Assume towards a contradiction that $\vec{\mathcal{W}}'$ is a proper initial segment of $\vec{\mathcal{W}}$, i.e. $\vec{\mathcal{W}}' \cap L_\vartheta[\vec{\mathcal{W}}] \cap L[\vec{\mathcal{W}}'] = \vec{\mathcal{W}} \upharpoonright \beta \cap L_\vartheta[\vec{\mathcal{W}}] \cap L[\vec{\mathcal{W}}']$ and there is a measure in $\vec{\mathcal{W}}$ above β . It easily follows that $L_\vartheta[\vec{\mathcal{W}}'] = L_\vartheta[\vec{\mathcal{W}} \upharpoonright \beta]$. The top measure in $\vec{\mathcal{W}}$ can be iterated making ϑ arbitrarily large, hence $L[\vec{\mathcal{W}}'] = L[\vec{\mathcal{W}} \upharpoonright \beta]$ and $\vec{\mathcal{W}}' = \vec{\mathcal{W}} \upharpoonright \beta \cap L[\vec{\mathcal{W}} \upharpoonright \beta]$. Moreover the iterated ultraprodut of $L_\vartheta[\vec{\mathcal{W}}]$ by the top measure produces a class of indiscernibles (defined in $L[\vec{\mathcal{U}}]$) for $L[\vec{\mathcal{W}}']$ containing all large

enough cardinals. That gives a definition in $L[\vec{\mathcal{U}}]$ of truth in $L[\vec{\mathcal{W}}']$: $L[\vec{\mathcal{W}}'] \models \sigma$ iff for all regular cardinals λ in $L[\vec{\mathcal{U}}]$ large enough $V_\lambda \cap L[\vec{\mathcal{W}}'] \models \sigma$. Consequently using the elementary embedding i' we obtain a definition of truth in $L[\vec{\mathcal{U}}]$: $L[\vec{\mathcal{U}}] \models \sigma$ iff for all regular cardinals λ large enough $V_\lambda \cap L[\vec{\mathcal{U}}] \models \sigma$. A contradiction. $\square$

Remark Notice that if $\vec{\mathcal{U}}$ is a coherent sequence of measures such that $o^\mathcal{U}(\kappa) < \kappa^{++}$ then the measures $\{U_\alpha^\kappa; \alpha < o^\mathcal{U}(\kappa)\}$ cannot cover $\mathcal{P}(\kappa^+)$: let $\gamma < \kappa^{++}$ be such that $\gamma > \kappa^{++\text{Ult}(V, U_\alpha^\kappa)}$ for all $\alpha < o^\mathcal{U}(\kappa)$, and let $A \subseteq \kappa^+$ code the well-ordering of order type γ . Then $A \notin \text{Ult}(V, U_\alpha^\kappa)$ for all $\alpha < o^\mathcal{U}(\kappa)$. Hence $o^\mathcal{U}(\kappa) = \kappa^{++}$ is a necessary condition for the covering of $\mathcal{P}(\kappa^+)$. However not sufficient: Let $\vec{\mathcal{U}}$ be a coherent sequence of measures in V , $o^\mathcal{U}(\kappa) = \kappa^{++}$. Let P be the Cohen forcing adding a subset of κ^+ , $G \subseteq \kappa^+$ P -generic over V . Since P is κ -closed no new κ -sequences are added, and so $\vec{\mathcal{U}}$ is also a coherent sequence of measures in $V[G]$. I claim that G is not covered by any of the measures $\{U_\alpha^\kappa; \alpha < \kappa^{++}\}$. Consider the ultraproduct embedding

$$j_\alpha^* : V[G] \rightarrow \text{Ult}(V[G], U_\alpha^\kappa) = M_\alpha[G^*]$$

which extends $j_\alpha : V \rightarrow \text{Ult}(V, U_\alpha^\kappa) = M_\alpha$. $G^* = j_\alpha^*(G)$ is $j_\alpha P$ -generic over M_α . But $j_\alpha P$ is the Cohen forcing adding a subset of $j_\alpha(\kappa^+)$ in the sense of M_α , so it is $j_\alpha \kappa$ -closed in M_α , in particular κ^+ -closed, and no new subsets of κ^+ are added by G^* . As $G \notin M_\alpha$ we conclude that $G \notin M_\alpha[G^*] = \text{Ult}(V[G], U_\alpha^\kappa)$. It can be actually proved using Mitchell's methods that if $V = L[\vec{\mathcal{U}}]$ then there are no new measures on κ in $V[G]$.

Now let us use the Laver's function on κ to construct closed systems of measures isomorphic to a given well-founded poset.

Proposition 3.5. *If there is a Laver's function on κ then there are two functions $F, G : \kappa \rightarrow V_\kappa$ such that*

$$\forall A, B \subseteq V_{\kappa+2}, |A| \leq \kappa^+, |B| \leq \kappa^+ \exists U \text{ a measure on } \kappa :$$

$$(j_U F)(\kappa) = A \text{ and } (j_U G)(\kappa) = B.$$

Proof. All we need is an effective coding of pairs (A, B) , where $A, B \subseteq V_{\lambda+2}$, $|A| \leq \lambda^+$, $|B| \leq \lambda^+$, by elements of $V_{\lambda+2}$. Firstly for $x, y \in V_{\lambda+2}$ define

$$x \oplus y = \{\{0\} \times z; z \in x\} \cup \{\{1\} \times z; z \in y\}.$$

It is easy to see that $x \oplus y \in V_{\lambda+2}$ if λ is a limit ordinal. On the other hand for any $z \in V_{\lambda+2}$ we can find unique $x, y \in V_{\lambda+2}$ such that $z = x \oplus y$ if there are any. Given $A = \{x_i; i \in I\} \subseteq V_{\lambda+2}$ indexed over a set $I \subseteq V_{\lambda+1}$ define

$$\bigoplus_{i \in I} x_i = \{i \times z; i \in I \text{ and } z \in x_i\}$$

which is in $V_{\lambda+2}$ if λ is a limit ordinal. If $z \in V_{\lambda+2}$ we can again find unique $I \subseteq V_{\lambda+1}$ and $A = \{x_i; i \in I\} \subseteq V_{\lambda+2}$ such that $z = \bigoplus_{i \in I} x_i$ if there are any. Finally code (A, B) , $A = \{x_i; i \in I\}$, $B = \{y_j; j \in J\}$ as

$$A \oplus B = \left(\bigoplus_{i \in I} x_i \right) \oplus \left(\bigoplus_{j \in J} y_j \right).$$

Let $f : \kappa \rightarrow V_\kappa$ be the Laver's function. For $\lambda < \kappa$ limit put $F(\lambda) = A_\lambda$, $G(\lambda) = B_\lambda$ if $f(\lambda) = A_\lambda \oplus B_\lambda$ for some A_λ, B_λ , otherwise $F(\lambda) = \emptyset = G(\lambda)$. For a given pair (A, B) find a measure U on κ such that $(j_U f)(\kappa) = A \oplus B$, then by the construction $(j_U F)(\kappa) = A$ and $(j_U G)(\kappa) = B$. $\square$

Proposition 3.6. *Let P be a well-founded poset of cardinality $\leq \kappa^+$. Then there are two closed systems of measures $S = \langle S_\lambda; \lambda \leq \kappa \rangle$, $T = \langle T_\lambda; \lambda \leq \kappa \rangle$ such that $S_\kappa = T_\kappa$, $\triangleleft_T$ extends $\triangleleft_S$, $P \cong (S_\kappa, \triangleleft_S)$ and T forms a coherent sequence of measures with $o_T(\kappa) \leq o(P) \cdot \kappa^+$. Moreover if U_0 is the first measure in T_κ then we can require that $j_{U_0}(\kappa)$ is greater than a given ordinal $\vartheta < \kappa^{++}$.*

Proof. Let $F, G : \kappa \rightarrow V_\kappa$ be the functions from 3.5. Enumerate $P = \{p_\alpha; \alpha < \mu\}$ so that $p_\alpha <_P p_\beta$ implies $\alpha < \beta$ and each level of P corresponds in this ordering to a segment of order type at most κ^+ . Consequently $\mu \leq o(P) \cdot \kappa^+$. Firstly find U_{p_0} such that $(j_{U_{p_0}} F)(\kappa) = \vartheta$ and $(j_{U_{p_0}} G)(\kappa) = \emptyset$. Then $j_{U_{p_0}}(\kappa) > \vartheta$ because $(j_{U_{p_0}} F)(\kappa) \in V_{j_{U_{p_0}}(\kappa)}$. Using the two functions F, G by induction on $\alpha > 0$ find U_{p_α} so that

$$(j_{U_{p_\alpha}} F)(\kappa) = \{U_{p_\gamma}; \gamma < \alpha\} \text{ and}$$

$$(j_{U_{p_\alpha}} G)(\kappa) = \{U_{p_\gamma}; p_\gamma <_P p_\alpha\}.$$

That gives $T_\kappa = S_\kappa = \{U_{p_\alpha}; \alpha < \mu\}$, put $T_\lambda = F(\lambda)$ and $S_\lambda = G(\lambda)$ if those are sets of measures over λ . Use lemma 3.1 to get closed systems of measures with required properties. $\square$

The proof works for well-founded κ^{++} -like posets P as well (i.e. $|P| \leq \kappa^{++}$ and $\forall p \in P : |P \restriction p| \leq \kappa^+$). We will need the auxiliary coherent sequence of measures T to make sure the measures in S_κ are separable.

The assumption $o^\mathcal{U}(\kappa) = \kappa^{++}$ can be significantly weakened to represent smaller well-founded posets. The proofs of 3.3, 3.6 can be modified to prove that if $o^\mathcal{U}(\kappa) = \kappa^+$ then all well-founded κ^+ -like posets P (i.e. $|P| \leq \kappa^+$, and $\forall p \in P : |P \restriction p| \leq \kappa$) can be represented as $\langle S_\kappa, \triangleleft_S \rangle$ for a closed system of measures S .

Separability and Repeat Points.

To prove theorem 2 let us give some estimates on the order of a least repeat point.

Fix a coherent sequence $\vec{U}$ with the least repeat point on κ of order Θ . We say that f_ν is an α -canonical function for ν if for all $\delta \in [\alpha, o^\mathcal{U}(\kappa)) : [f_\nu]_{U_\delta^\kappa} = \nu$.

Lemma 3.7. [JW93] *If U_α^κ is not a repeat point then α has an α -canonical function. Moreover there is $X_\alpha \subseteq \kappa$ such that*

$$\forall \delta < o^\mathcal{U}(\kappa) : X_\alpha \in U_\alpha^\kappa \leftrightarrow \delta = \alpha.$$

Proof. Let $X \in U_\alpha^\kappa$ be such that $X \notin U_\nu^\kappa$ for $\nu < \alpha$. Put

$$f_\alpha(\xi) = \sup\{\eta \leq o^\mathcal{U}(\xi); \forall \eta' < \eta : X \cap \xi \notin U_{\eta'}^\kappa\}.$$

Let $\delta \geq \alpha$, $j : V \rightarrow M = \text{Ult}(V, U_\delta^\kappa)$ then

$$M \models (jf_\alpha)(\kappa) = \sup\{\eta \leq o^{j\mathcal{U}}(\kappa) = \delta; \forall \eta' < \eta : X \notin U_{\eta'}^\kappa\}.$$

Hence $(jf_\alpha)(\kappa) = \alpha$ since $X \in U_\alpha^\kappa$ and $X \notin U_{\eta'}^\kappa$ for $\eta' < \alpha$. Put $Y = \{\xi < \kappa; f_\alpha(\xi) \geq o^\mathcal{U}(\xi)\}$, then $Y \in U_\delta^\kappa$ iff $\delta \leq \alpha$. Consequently $X \cap Y$ separates U_α^κ from all the other measures. $\square$

It means that if there are no repeat points on κ in $\vec{\mathcal{U}}$ then the measures on κ are separable.

Lemma 3.8. *If $\nu > \alpha$ has an α -canonical function and α is not a repeat point then ν is not a repeat point.*

Proof. Let f_α be the α -canonical function for α defined above, and f_ν an α -canonical function for ν . Put $A = \{\xi < \kappa; o^\mathcal{U}(\xi) > f_\alpha(\xi)\}$,

$$B = A \cap \{\xi < \kappa; o^\mathcal{U}(\xi) = f_\nu(\xi)\}.$$

Then $A \in U_\delta^\kappa$ iff $\delta > \alpha$, and so $B \in U_\nu$ separates U_ν from all the other measures. $\square$

Consequently if α is not a repeat point then $\alpha + 1$ is not a repeat point as $f_{\alpha+1}(\xi) = f_\alpha(\xi) + 1$ is an α -canonical function for $\alpha + 1$. The following is a joint result with J.Zapletal.

Proposition 3.9. *If $\alpha < \Theta$ then $j_{U_\alpha^\kappa}(\kappa) < \Theta$.*

Proof. Put $\gamma = j_{U_\alpha^\kappa}(\kappa)$, $\alpha < \Theta$. Let f_α be the α -canonical function for α . We prove that for any $\nu < \gamma$, U_ν is not a repeat point. Let $g_\nu : \kappa \rightarrow \kappa$ be a function such that $(j_{U_\alpha^\kappa} g_\nu)(\kappa) = \nu$. Define for $\lambda < \kappa$

$$\tilde{g}_\nu(\lambda) = \begin{cases} (j_{U_{f_\alpha(\lambda)}^\kappa}^\lambda(g_\nu \upharpoonright \lambda))(\lambda) & \text{if } o^\mathcal{U}(\lambda) > f_\alpha(\lambda) \\ g_\nu(\lambda) & \text{otherwise .} \end{cases}$$

I claim that $\tilde{g}_\nu$ is an α -canonical function for ν . By the choice of g_ν : $(j_{U_\alpha^\kappa} \tilde{g}_\nu)(\kappa) = \nu$. Let $\delta \in (\alpha, o^\mathcal{U}(\kappa))$ then

$$(j_{U_\delta^\kappa} \tilde{g}_\nu)(\kappa) = (j_{U_\alpha^\kappa}^{M_\delta}((j_{U_\delta^\kappa} g_\nu) \upharpoonright \kappa))(\kappa) = (j_{U_\alpha^\kappa} g_\nu)(\kappa) = \nu$$

since $(j_{U_\delta^\kappa} g_\nu) \upharpoonright \kappa = g_\nu$ and $j_{U_\alpha^\kappa}^{M_\delta} = j_{U_\alpha^\kappa} \upharpoonright M_\delta$, where $M_\delta = \text{Ult}(V, U_\delta^\kappa)$. According to 3.8 U_ν^κ is not a repeat point.

Consequently $\gamma \leq \Theta$. Note that in general $U \triangleleft W$ implies that $j_U(\kappa) < j_W(\kappa)$. Since $\alpha + 1 < \Theta$ we see that $\Theta \geq j_{U_{\alpha+1}^\kappa}(\kappa) > j_{U_\alpha^\kappa}(\kappa)$. $\square$

Remark. Θ is a limit ordinal between κ^+ and κ^{++} of cofinality κ^+ . Moreover if F is a Σ_1 ordinal operation and $\bar{\alpha} < \Theta$ then $F(\bar{\alpha}) < \Theta$.

Proof. Let F be given by a Σ_1 formula $\varphi(\bar{\alpha}, \beta)$ such that $ZFC \vdash \text{“}\varphi(\bar{\alpha}, \beta) \text{ is a function of } \bar{\alpha} \text{”}$. Let $\bar{\alpha} < \Theta$, find a $\delta < \Theta$ such that $\bar{\alpha} < j_{U_\delta}(\kappa) = \tilde{\kappa}$ ($\delta > \bar{\alpha}$ is sufficient). Since $M_\delta \models \text{“}\tilde{\kappa} \text{ is inaccessible,”}$ i.e. $V_{\tilde{\kappa}}^{M_\delta} \models ZFC$, it is necessary that

$$M_\delta \models \forall \bar{\beta} < \tilde{\kappa} : F^{V_{\tilde{\kappa}}}(\bar{\beta}) < \tilde{\kappa}.$$

Moreover $M_\delta \models \text{“}\forall \bar{\beta} < \tilde{\kappa} : F^{V_{\tilde{\kappa}}}(\bar{\beta}) = F(\bar{\beta}) \text{”}$ and $F^{M_\delta} = F^V$ as F is a Σ_1 function. Hence $F(\bar{\alpha}) < \tilde{\kappa} < \Theta$. $\square$

In particular Θ is inaccessible by primitive recursive ordinal operations.

Finally we are ready to prove theorem 2.

Proof of theorem 2. Given a well-founded poset P enumerate P as in the proof of 3.6, $P = \{p_\alpha; \alpha < \mu\}$, $\mu < \kappa^{++}$. Then use 3.6 to find a closed system of measures $\langle S_\lambda; \lambda \leq \kappa \rangle$ such that $P \cong (S_\kappa, \triangleleft_S)$, and a coherent sequence $\langle T_\lambda; \lambda \leq \kappa \rangle$ such that $T_\kappa = S_\kappa$ and the first measure U_0 satisfies $j_{U_0}(\kappa) > \mu$. Then by 3.9 T_κ cannot have a repeat point as $o_T(\kappa) < j_{U_0}(\kappa)$. Consequently the measures in $S_\kappa = T_\kappa$ are separable. $\square$

REFERENCES

- [J84] T. Jech, *Stationary subsets of inaccessible cardinals*, Contemporary Mathematics **31** (1984), 115–141.
- [J86] T. Jech, *Multiple Forcing*, Cambridge University Press, 1986.
- [J89] T. Jech, *Positive Σ operations on cardinals and normal filters on greatly Mahlo cardinals*, JSL **54** (1989), 226–233.
- [JS90] T. Jech, S. Shelah, *Full reflection of stationary sets below $\aleph_\omega$* , JSL **55** (1990), 822–829.
- [JS93] T. Jech, S. Shelah, *Full reflection of stationary sets at regular cardinals*, American Journal of Mathematics **115** (1993), 435–453.
- [JW93] T. Jech, J. Witzany, *Full reflection at a measurable cardinal*, JSL, to appear in JSL.
- [Ka93] A. Kanamori, *The Higher Infinite I, II*, in preparation.
- [La78] R. Laver, *Making supercompactness of κ indestructible under κ -directed closed forcing*, Israel J. Math **29** (1978), 385–388.
- [M82] M. Magidor, *Reflecting stationary sets*, JSL **47** (1982), 755–771.
- [Mi83] W. J. Mitchell, *Sets constructible from sequences of measures: revisited*, JSL **48** (1983), 600–609.
- [WoC92] H. Woodin, J. Cummings, *Generalised Prikry Forcings*, in preparation.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
PA 16802

E-mail address: witzany@math.psu.edu